








# Can additive beta diversity be reliably partitioned into nestedness and turnover components?

Thomas J. Matthews<sup>1,2,3</sup>  | Thomas W. H. Aspin<sup>1</sup> | Werner Ulrich<sup>4</sup>  |  
 Andrés Baselga<sup>5</sup> | Yasuhiro Kubota<sup>6</sup>  | Konstantinos Proios<sup>7</sup>  | Kostas A. Triantis<sup>7</sup>  |  
 Robert J. Whittaker<sup>8,9</sup>  | Giovanni Strona<sup>10</sup> 

<sup>1</sup>GEES (School of Geography, Earth and Environmental Sciences), The University of Birmingham, Birmingham, United Kingdom

<sup>2</sup>CE3C – Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity Group and Univ. dos Açores – Depto de Ciências e Engenharia do Ambiente, Açores, Portugal

<sup>3</sup>Birmingham Institute of Forest Research, University of Birmingham, Birmingham, United Kingdom

<sup>4</sup>Faculty of Biology and Environment Protection, Nicolaus Copernicus University, Toruń, Poland

<sup>5</sup>Facultad de Biología, Departamento de Zoología, Genética y Antropología Física, Universidad de Santiago de Compostela, Santiago de Compostela, Spain

<sup>6</sup>Faculty of Science, University of the Ryukyus, Nishihara, Japan

<sup>7</sup>Faculty of Biology, Department of Ecology and Taxonomy, National and Kapodistrian University of Athens, Athens, Greece

<sup>8</sup>School of Geography and the Environment, University of Oxford, Oxford, United Kingdom

<sup>9</sup>Center for Macroecology, Evolution and Climate, National Museum of Natural History, University of Copenhagen, Copenhagen, Denmark

<sup>10</sup>European Commission, Joint Research Centre, Directorate D – Sustainable Resources, Ispra (VA), Italy

## Correspondence

Thomas J. Matthews, GEES (School of Geography, Earth and Environmental Sciences), The University of Birmingham, Birmingham B15 2TT, UK.  
 Email: t.j.matthews@bham.ac.uk

Editor: Petr Keil

## Abstract

**Aims:** Quantifying  $\beta$ -diversity (differences in the composition of communities) is central to many ecological studies. There are many  $\beta$ -diversity metrics, falling mostly into two approaches: variance-based (e.g., the Sørensen index), or diversity partitioning (e.g., additive  $\beta$ -diversity). The former cannot be used when species–sites matrices are unavailable (which is often the case in island biogeography in particular) and only species richness data are provided. Recently, efforts have been made to partition additive  $\beta$ -diversity, a metric calculated using only  $\alpha$ -diversity and  $\gamma$ -diversity, into nestedness and turnover components (termed here “richness-only  $\beta$ -diversity partitioning”). We set out to test whether this form of  $\beta$ -diversity partitioning generates interpretable results, comparable with metrics based on species incidence  $\beta$ -diversity partitioning.

**Location:** Global.

**Time period:** Present day.

**Major taxa studied:** Multiple taxa.

**Methods:** We first provide a brief review of  $\beta$ -diversity partitioning methods, with a particular focus on the development of richness-only  $\beta$ -diversity partitioning. Second, we use 254 empirical incidence matrices (provided with the paper) sourced from the literature to measure turnover and nestedness using incidence  $\beta$ -diversity partitioning, comparing the resulting values with those calculated using richness-only  $\beta$ -diversity.

**Results:** We provide an account of the emergence of  $\beta$ -diversity partitioning, with particular reference to the analysis of richness-only datasets, and to the definition and usage of the relevant metrics. Analytically, we report weak correlations between turnover and nestedness calculated using the two different approaches. We show that this is because identical values of  $\alpha$ -diversity and  $\gamma$ -diversity can correspond to incidence matrices with a range of different structures.

**Main conclusions:** Our results demonstrate that the use of richness-only  $\beta$ -diversity partitioning to measure turnover and nestedness is problematic and can produce patterns unrelated to conventional measures of turnover and nestedness. We

therefore recommend that more accurate definitions are adopted for these terms in future studies.

#### KEYWORDS

beta diversity, compositional differences, diversity partitioning, nestedness, turnover

## 1 | INTRODUCTION

Quantifying differences in the composition of communities (i.e., measuring  $\beta$ -diversity) and testing the prevalence of nestedness in ecological communities are central to many ecological studies (Rosenzweig, 1995; Ulrich, Almeida-Neto, & Gotelli, 2009; Whittaker, 1960). Various metrics have been proposed to measure  $\beta$ -diversity (Anderson et al., 2011; Chao, Chiu, & Hsieh, 2012; Koleff, Gaston, & Lennon, 2003; Tuomisto, 2010), and many of these can be broadly divided into variance-based approaches (e.g., the Sørensen and Jaccard indices) and diversity partitioning-based approaches (Chao & Chiu, 2016; Legendre & De Cáceres, 2013). Recent work has bridged these two approaches to calculating  $\beta$ -diversity (Chao & Chiu, 2016; Chao et al., 2012). Nestedness, as originally conceived within island biogeography, refers to the ordered loss/gain of species along a richness gradient, whereby each larger assemblage tends to contain all the members of the previously considered site (Matthews, Cottee-Jones, & Whittaker, 2015; Patterson & Atmar, 1986; Ulrich et al., 2009). The calculation of most nestedness and variance-based  $\beta$ -diversity metrics requires knowledge of species occurrences at each site, coded in a binary presence-absence matrix (herein, incidence matrix), with species in rows and sites in columns (e.g., Koleff et al., 2003; Legendre & Legendre, 1983; Ulrich et al., 2009). Table 1 provides a glossary of the (many) different metrics discussed in the present study, and Figure 1 provides an illustration of the different metrics.

### 1.1 | Diversity partitioning and richness-only $\beta$ -diversity partitioning: An overview

Over time, as research questions and meta-analytical tools have developed, the published literature has become an increasingly important source of data to extend the power of analyses via data mining. This is particularly true in regard to studies that conduct meta-analyses of  $\beta$ -diversity and nestedness (e.g., Cabral, Weigelt, Kissling, & Kreft, 2014; Matthews et al., 2015). However, in many cases the full incidence matrix for a set of sites is not available in previously published studies, and only the number of species in each site ( $\alpha$ -diversity) and the regional diversity ( $\gamma$ -diversity) can be retrieved (e.g., Cowie, 1995). This is a particular problem in island biogeography, where several meta-analyses of ecological patterns on islands are based on these simple data [see, for example, various studies investigating species-area relationships (SARs), such as Matthews, Guilhaumon, Triantis, Borregaard, & Whittaker, 2016; Triantis, Guilhaumon, & Whittaker, 2012]. Authors have attempted

to get around this problem by estimating  $\beta$ -diversity and nestedness through diversity partitioning approaches (e.g., Cabral et al., 2014; Chiarucci, Bacaro, Arévalo, Delgado, & Fernández-Palacios, 2010; Sfenthourakis & Panitsa, 2012; see Zhang et al., 2014, for a terrestrial example).

Estimating  $\beta$ -diversity using diversity partitioning has a long history in ecology (Jost, 2007; Lande, 1996; MacArthur, Recher, & Cody, 1966; Tuomisto, 2010; Whittaker, 1960, 1965). The two main diversity partitioning approaches used are additive ( $\beta_{\text{Total}} = \gamma - \alpha$ ), where  $\beta_{\text{Total}}$  is the amount by which regional diversity ( $\gamma$ ) exceeds the mean diversity of a set of sites ( $\alpha$ ), and multiplicative ( $\beta_{\text{Mult}} = \gamma/\alpha$ ), where  $\beta_{\text{Mult}}$  is the regional-to-local diversity ratio (Crist, Veech, Gering, & Summerville, 2003; Tuomisto, 2010; Veech, Summerville, Crist, & Gering, 2002; Whittaker, 1960). Their relative merits have been much debated (see Baselga, 2010a; Crist et al., 2003; Jost, 2007; Lande, 1996; Tuomisto, 2010; Veech & Crist, 2010), and recent work has shown how they (i.e., additive and multiplicative diversity partitioning) are mathematically linked (Chao et al., 2012). Additively partitioned  $\beta$ -diversity ( $\beta_{\text{Total}}$ ), which is the focus of this present study, has been shown to be dependent on both  $\gamma$  and the number of sites ( $N$ ), and it has thus been argued that  $\beta_{\text{Total}}$  should be normalized (by  $\gamma$  and  $N$ ;  $\beta_{\text{Stan}}$ ) in order to compare  $\beta$ -diversity values (Chao et al., 2012).

Using an additive partitioning approach,  $\beta_{\text{Total}}$  has been further partitioned into two sub-components, which have been argued to measure nestedness and turnover (we term this “richness-only  $\beta$ -diversity partitioning”). This is different from the partitioning of dissimilarity indices (such as Sørensen and Jaccard indices), through analysis of an incidence matrix (we term this “incidence  $\beta$ -diversity partitioning”), into turnover and nestedness-resultant dissimilarity/richness difference components (Baselga, 2010b, 2012; Carvalho, Cardoso, & Gomes, 2012). The use of richness-only  $\beta$ -diversity partitioning appears to have been based on the approach of Crist and Veech (2006), who used the power law SAR model to partition  $\beta_{\text{Total}}$  in isolated habitats into two components:  $\beta_{\text{Area}}$ , which is intended to describe how much of  $\beta_{\text{Total}}$  is due to the area effect, and  $\beta_{\text{Replace}}$ , which is intended to describe how much is explained by other factors (see Crist & Veech, 2006).  $\beta_{\text{Area}}$  is defined by Crist and Veech (2006, p. 928) as “the mean deviation between the species richness of the largest habitat patch and the species richness of smaller patches”. Subsequent studies have used the  $\beta_{\text{Area}}$  component as a measure of nestedness, and the  $\beta_{\text{Replace}}$  component as a measure of replacement/turnover (e.g., Cabral et al., 2014; Chiarucci et al., 2010; Sfenthourakis & Panitsa, 2012; Zhang et al., 2014). It is important to note that the interpretation of  $\beta_{\text{Area}}$  and  $\beta_{\text{Replace}}$  as measures of

Metric	Description	References
<i>Incidence matrix metrics</i>		
$\beta_{\text{Sor}}$	Sørensen dissimilarity; a $\beta$ -diversity metric that measures compositional differences between sites	Baselga (2010b, 2012)
$\beta_{\text{Sim}}$	Simpson dissimilarity; the turnover component of Sørensen dissimilarity	Baselga (2010b, 2012)
$\beta_{\text{Sne}}$	Nestedness-resultant fraction of Sørensen dissimilarity	Baselga (2010b, 2012)
NODF	A nestedness index based on the twin properties of standardized differences in matrix row and column fills and paired overlap	Almeida-Neto et al. (2008)
<i>Richness-only metrics</i>		
$\beta_{\text{Total}}$	$\beta$ -diversity calculated using additive diversity partitioning; the amount that regional diversity exceeds the mean diversity of a set of sites	MacArthur et al. (1966), Lande (1996), Veech et al. (2002)
$\beta_{\text{Stan}}$	$\beta_{\text{Total}}$ normalized by $\gamma$ and the number of sites	Chao et al. (2012)
$\beta_{\text{Nestedness}}$	Hitherto interpreted as a measure of nestedness. Its calculation is almost identical to the $\beta_{\text{Area}}$ metric of Crist and Veech (2006) but uses the maximum observed richness of an island in the dataset rather than maximum richness predicted by the power law species–area relationship model	Chiarucci et al. (2010), Cabral et al. (2014), Zhang et al. (2014)
$\beta_{\text{Replacement}}$	The turnover component of $\beta_{\text{Total}}$ . Hitherto interpreted as a measure of the compositional differences across a set of sites. Similar in calculation to the $\beta_{\text{Replace}}$ of Crist and Veech (2006)	Chiarucci et al. (2010), Cabral et al. (2014), Zhang et al. (2014)
$\beta_{\text{Area}}$	Measures the portion of $\beta_{\text{Total}}$ that is due to area effects	Crist and Veech (2006)
$\beta_{\text{Replace}}$	The portion of $\beta_{\text{Total}}$ that is due to factors other than area	Crist and Veech (2006)
$\beta_{\text{Mult}}$	$\beta$ -diversity calculated using multiplicative diversity partitioning; the regional-to-local diversity ratio (true $\beta$ -diversity)	Whittaker (1960, 1965), Jost (2007), Tuomisto (2010)

Note: The metrics are split into those that are calculated using the incidence matrix and those that are calculated using simple richness data (i.e.,  $\gamma$ - and  $\alpha$ -diversity).


nestedness and turnover, respectively, was not necessarily implied in the original study (Crist & Veech, 2006), a fact recognized by at least one of the subsequent studies (Cabral et al., 2014).

To take one study that used richness-only  $\beta$ -diversity partitioning as an example, Chiarucci et al. (2010, p. 86), in their study of plants on a variety of Macaronesian islands, use the term  $\beta_{\text{Nestedness}}$  (rather than  $\beta_{\text{Area}}$ ), stating that “the first component of [additive]  $\beta$ -diversity ( $\beta_{\text{Nestedness}}$ ) quantified the degree of nestedness of the flora,” whilst the “second  $\beta$  component ( $\beta_{\text{Replacement}}$ ) measure[s] the differences in species composition among the flora of the islands within an archipelago, and [is] a measure of the compositional differences across islands.” The authors then proceed to make inferences regarding the nestedness of their data; for example, “the higher importance of  $\beta_{\text{Nestedness}}$  for pteridophytes indicated that, for this taxon, the flora of each island is largely formed by a subset of species that make up the archipelago flora” (Chiarucci et al., 2010, p. 89). This example, and others (e.g., Cabral et al., 2014; Sfenthourakis & Panitsa, 2012; Zhang et al., 2014), illustrates that colleagues have started to use richness-only

**TABLE 1** A glossary of the different  $\beta$ -diversity and nestedness metrics used in this study

$\beta$ -diversity partitioning in their research; the use of the method and its implications is not confined to those four cited studies and a simple Google Scholar search indicates these four papers have been cited over 80 times. It is therefore timely to assess the implications of this approach and how well the richness-only  $\beta$ -diversity partitions correspond with conventional measures of nestedness and turnover, as this could constitute a useful analytical tool if it can be shown to be robust.

In this article, we use a dataset of 254 incidence matrices (details below) to assess to what extent nestedness and turnover calculated by richness-only  $\beta$ -diversity partitioning (i.e., partitioning additive  $\beta$ -diversity, i.e.,  $\beta_{\text{Total}}$ ) are congruent with nestedness and compositional difference metrics calculated using the full incidence matrix. It is important to re-stress that we are focused on the issues surrounding the use of richness-only  $\beta$ -diversity partitioning (i.e., when there is no incidence matrix and thus no information on which species are present on which island; e.g., as employed by Chiarucci et al., 2010), and not incidence  $\beta$ -diversity partitioning (e.g., the partitioning of Sørensen dissimilarity; e.g., Baselga, 2012). As  $\beta_{\text{Total}}$  is known to be



**FIGURE 1** Overview of the various total  $\beta$ -diversity, turnover and nestedness metrics discussed in the main text. The metrics are organized within the figure according to how they are calculated (richness-only  $\beta$ -diversity partitioning and incidence matrix analysis), and to what it is they are purported to be measuring (total  $\beta$ -diversity, turnover or nestedness). The two solid black boxes indicate metrics that have been partitioned into two components (connected by the black lines). Here, richness-only  $\beta$ -diversity partitioning relates to the partitioning of additive  $\beta$ -diversity. For definitions of terms, see Table 1

problematic due to its dependency on both  $\gamma$  and the number of sites ( $N$ ), we might expect the partitioned components of  $\beta_{\text{Total}}$  to also have issues. However, this possibility has not previously been explored, and it is important that any problems with the approach are highlighted to avoid the proliferation of incorrect metrics (or at least the incorrect interpretation of particular metrics) in the ecological literature.

## 2 | MATERIALS AND METHODS

To compare nestedness and turnover calculated using richness-only  $\beta$ -diversity partitioning (i.e., additive partitioning of  $\beta_{\text{Total}}$ ) with nestedness and compositional difference metrics based on incidence  $\beta$ -diversity partitioning (i.e., analysis of the full incidence matrix), we used a collection of island incidence matrices that we sourced from the literature. Briefly, the database contains 254 incidence matrices of various taxa from different island systems (each representing a geographically coherent set of islands), including all major island types such as volcanic oceanic islands, continental-shelf islands, atolls, and habitat islands. Unlike other collections of incidence matrices (e.g., Atmar & Patterson, 1995; Strona, Ulrich, & Gotelli, 2017), our database is comprised solely of island datasets and thus allows us to make general conclusions regarding patterns of interest in islands (Matthews, 2015; Whittaker & Fernández-Palacios, 2007). The full database will be published as part of a separate upcoming study, but the set of 254 incidence matrices are available from GitHub (txm676/Partitioning-additive-beta). The repository has been archived on the Zenodo research data repository (<https://doi.org/10.5281/zenodo.2595322>).

For each of the 254 datasets, we calculated additive  $\beta$ -diversity ( $\beta_{\text{Total}}$ ). Thus, the incidence matrices were not used; we simply used

mean  $\alpha$  (the average richness of the set of islands in a dataset) and  $\gamma$  (the overall species richness of the set of islands in a dataset). We also calculated normalized  $\beta$ -diversity ( $\beta_{\text{Stan}}$ ) using the approach in Chao et al. (2012):

$$\beta_{\text{Stan}} = \frac{\beta_{\text{Total}}}{\left(1 - \frac{1}{N}\right)\gamma}, \quad (1)$$

where  $N$  is the number of islands. We then calculated the  $\beta_{\text{Nestedness}}$  and  $\beta_{\text{Replacement}}$  (Table 1) partitions of  $\beta_{\text{Total}}$  (i.e., richness-only  $\beta$ -diversity partitioning) using the approach of Chiarucci et al. (2010). This approach differs slightly from that employed by Crist and Veech (2006) in that it uses the maximum observed richness of an island in the dataset rather than maximum richness predicted by the power law SAR model. However, the two approaches produce highly correlated values [Spearman's rho = .88,  $p < .001$ , for the correlation between the  $\beta_{\text{Nestedness}}$  of Chiarucci et al. (2010) and the  $\beta_{\text{Area}}$  of Crist and Veech (2006) based on the 254 empirical matrices]. Following Chiarucci et al. (2010)  $\beta_{\text{Nestedness}}$  was calculated using the equation:

$$\beta_{\text{Nestedness}} = \frac{1}{N} \sum_{i=1}^N (S_{\text{max}} - S_i), \quad (2)$$

where  $S_i$  is the number of species on the  $i$ -th island, and  $S_{\text{max}}$  is the number of species on the most species rich island.  $\beta_{\text{Replacement}}$  can then be calculated using the equation:

$$\beta_{\text{Replacement}} = \beta_{\text{Total}} - \beta_{\text{Nestedness}}. \quad (3)$$

We note that Equation 2 can be reformulated to clarify the meaning of  $\beta_{\text{Nestedness}}$ :

$$\beta_{\text{Nestedness}} = \frac{1}{N} \sum_{i=1}^N (S_{\text{max}} - S_i) = S_{\text{max}} - \frac{1}{N} \sum_{i=1}^N S_i = S_{\text{max}} - \alpha. \quad (4)$$

This is in accordance with the initial definition of  $\beta_{\text{Area}}$  by Crist and Veech (2006). As the average local diversity,  $\alpha$ , is independent of the number of sites considered  $\beta_{\text{Nestedness}}$  can be derived from the knowledge of two sites having  $S_{\text{max}}$  and  $S_r$  species,

$$\beta_{\text{Nestedness}} = S_{\text{max}} - \frac{S_{\text{max}} + S_r}{2} = \frac{S_{\text{max}} - S_r}{2}, \quad (5)$$

Therefore,  $\beta_{\text{Nestedness}}$  reduces to a difference in species richness between two sites, irrespective of the occurrence of joint species among these sites. This contradicts the basic definition of nestedness, which asserts the existence of an ordered set of subsamples (Almeida-Neto, Guimarães, Guimarães, Loyola, & Ulrich, 2008; Patterson & Atmar, 1986). Taking the extreme case of there being no shared species, and thus with the maximum possible species turnover,  $\beta_{\text{Nestedness}}$  can take any of the full range of possible values between zero and  $S_{\text{max}} - \alpha$ .

We also measured the compositional difference between islands in a dataset using the Sørensen dissimilarity index computed on the full incidence matrix, using the “betapart” R package (version 1.4-1, Baselga & Orme, 2012). Overall compositional difference was calculated using Sørensen multi-site dissimilarity ( $\beta_{\text{Sor}}$ ; see Table 1),

$$\beta_{\text{Sor}} = \frac{\left[ \sum_{i<j} \min(b_{ij}, b_{ji}) \right] + \left[ \sum_{i<j} \max(b_{ij}, b_{ji}) \right]}{2 \left[ \sum_i S_i - S_T \right] + \left[ \sum_{i<j} \min(b_{ij}, b_{ji}) \right] + \left[ \sum_{i<j} \max(b_{ij}, b_{ji}) \right]} \quad (6)$$

where  $S_i$  is the total number of species in site  $i$ ,  $S_T$  is the total number of species in all sites considered together and  $b_{ij}$ ,  $b_{ji}$  are the number of species exclusive to sites  $i$  and  $j$ , respectively. Sørensen multi-site dissimilarity was partitioned (i.e., incidence  $\beta$ -diversity partitioning) into the turnover component (Simpson multi-site dissimilarity;  $\beta_{\text{Sim}}$ ; see Table 1),

$$\beta_{\text{Sim}} = \frac{\left[ \sum_{i<j} \min(b_{ij}, b_{ji}) \right]}{\left[ \sum_i S_i - S_T \right] + \left[ \sum_{i<j} \min(b_{ij}, b_{ji}) \right]} \quad (7)$$

and the nestedness component (Nestedness-resultant multi-site dissimilarity;  $\beta_{\text{Sne}}$ , Baselga, 2010b, 2012),

$$\beta_{\text{Sne}} = \frac{\left[ \sum_{i<j} \max(b_{ij}, b_{ji}) \right] - \left[ \sum_{i<j} \min(b_{ij}, b_{ji}) \right]}{2 \left[ \sum_i S_i - S_T \right] + \left[ \sum_{i<j} \min(b_{ij}, b_{ji}) \right] + \left[ \sum_{i<j} \max(b_{ij}, b_{ji}) \right]} \times \frac{\sum_i S_i - S_T}{\left[ \sum_i S_i - S_T \right] + \left[ \sum_{i<j} \min(b_{ij}, b_{ji}) \right]} \quad (8)$$


As  $\beta_{\text{Sne}}$  is conceptually distinct from “true” nestedness (Almeida-Neto, Frensel, & Ulrich, 2012; Baselga, 2012), we also measured the nestedness of each dataset with the nestedness metric based

on overlap and decreasing fill (NODF) metric (Almeida-Neto et al., 2008). According to NODF, an incidence matrix sorted in decreasing order of marginal totals is maximally nested when there is complete overlap of presence values (1s) from the right to the left column and from the bottom to the top row, and no ties in both the row and column marginal totals. We chose NODF as it is widely considered to be the most appropriate measure of nestedness (Ulrich et al., 2009). NODF was computed on the full incidence matrix, using the vegan R package (version 2.4-5, Oksanen et al., 2017). We used a combination of graphical plots and Kendall rank correlation tests to assess the correlation between the richness-only  $\beta$ -diversity partitioning metrics,  $\beta_{\text{Nestedness}}$  and  $\beta_{\text{Replacement}}$ , and the incidence  $\beta$ -diversity partitioning metrics,  $\beta_{\text{Sim}}$  and  $\beta_{\text{Sne}}$ , and also NODF. We used the Kendall rank correlation test rather than Spearman rank correlations as the former is argued to provide better results with small sample sizes. To illustrate that the issues we highlight lie with richness-only  $\beta$ -diversity partitioning and not with additive  $\beta$ -diversity ( $\beta_{\text{Total}}$ ) itself, we also assessed the correlation between  $\beta_{\text{Stan}}$  and  $\beta_{\text{Sor}}$  using a Kendall rank correlation test.

To determine whether our results were consistent across different types of island datasets, we divided our main dataset a number of different ways. First, we split the datasets into true islands (oceanic and continental-shelf islands, and islands within natural lakes) and habitat islands (all other datasets). Second, we calculated the quantiles, using all datasets, of each of: the number of islands,  $\alpha$ , and  $\gamma$ . For each factor, we then took the datasets representing the top and bottom quantiles (lowest and highest 25%) to be individual subsets. We then repeated the above analyses using each individual subset. Finally, we repeated the main correlation tests using the multi-site version of Jaccard dissimilarity calculated using both the Baselga (2012) and the Carvalho et al. (2012) approaches. The above analyses were undertaken using the R programming language (version 3.5.2, R Core Team, 2017), and the R code used to run the analyses is provided in a GitHub repository (txm676/Partitioning-additive-beta).

### 3 | RESULTS

Considering the full set of 254 empirical island incidence matrices, we found low (often very low) correlation between the relevant metrics. The lowest correlation was between  $\beta_{\text{Nestedness}}$  and  $\beta_{\text{Sne}}$  (Kendall's tau =  $-.04$ ;  $p$ -value =  $.36$ ), followed by  $\beta_{\text{Nestedness}}$  and NODF (Kendall's tau =  $-.11$ ;  $p$ -value =  $.01$ ),  $\beta_{\text{Replacement}}$  and  $\beta_{\text{Sor}}$  (Kendall's tau =  $.23$ ;  $p$ -value <  $.001$ ) and  $\beta_{\text{Replacement}}$  and  $\beta_{\text{Sim}}$  (Kendall's tau =  $.43$ ;  $p$ -value <  $.001$ ). As expected, there was a higher correlation between  $\beta_{\text{Stan}}$  and  $\beta_{\text{Sor}}$  (Kendall's tau =  $.56$ ;  $p$ -value <  $.001$ ). The plots of these relationships are provided in Figure 2; certain variables were logged prior to plotting (but not analysis) to ease the visual interpretation (see the legend of Figure 2). A complete set of pairwise scatter plots (along with their correlations) for all variables is provided as Figure S1 in Supporting Information Appendix S1.




**FIGURE 2** Scatter plots showing the relationship between different  $\beta$ -diversity and nestedness metrics, calculated using richness-only  $\beta$ -diversity partitioning and through analysing the full incidence matrix. The data are 254 empirical incidence matrices from different island systems (e.g., oceanic islands, habitat islands). The metrics on the x axis in all three plots are those calculated using richness-only  $\beta$ -diversity partitioning, whereas those on the y axis were calculated using the full incidence matrix. For the definitions of the variables, see Table 1.  $\beta_{\text{Nestedness}}$  and  $\beta_{\text{Replacement}}$  have been logged (base-e; a constant of 0.1 was added to all values to avoid zero values) for presentation purposes

Re-running the analyses using the different dataset subsets generated broadly similar results (Table S1 in Supporting Information Appendix S1). The main difference was for the correlation between  $\beta_{\text{Nestedness}}$  and  $\beta_{\text{Sne}}$ , whereby for three subsets ( $\alpha > 25$ ,  $\gamma < 20$  and  $\gamma > 77$ ) the correlation was positive and significant, although the correlation coefficient was less than .50 in all three cases (Supporting Information Table S1). The results of the analyses using the multi-site version of Jaccard dissimilarity calculated using both the Baselga (2012) and the Carvalho et al. (2012) approaches were similar to the main results and are not discussed further (Supporting Information Appendix S1).

## 4 | DISCUSSION

We have explored to what extent nestedness and turnover calculated through richness-only  $\beta$ -diversity partitioning (i.e., partitioning  $\beta_{\text{Total}}$ ) are congruent with nestedness and compositional difference metrics calculated using incidence  $\beta$ -diversity partitioning. We find that neither partition of  $\beta_{\text{Total}}$  provides good measures of what is commonly regarded as nestedness or turnover. Our results lead us to caution against the interpretation of the additive sub-components of  $\beta_{\text{Total}}$  as metrics of nestedness and replacement/turnover. The problem is most acute when considering nestedness:  $\beta_{\text{Nestedness}}$  and NODF calculated using the empirical matrices were in fact weakly negatively correlated (tau = -.11). A thought experiment that further illustrates the issue with  $\beta_{\text{Nestedness}}$  is provided in Supporting Information Appendix S2. Thus,  $\beta_{\text{Nestedness}}$  should simply be interpreted as representing area effects, as originally proposed by Crist and Veech (2006). Based on analyses of the empirical matrices, the correlation between  $\beta_{\text{Replacement}}$  and  $\beta_{\text{Sim}}$  was also low (tau = .43) and we do not recommend using  $\beta_{\text{Replacement}}$  in future studies to measure turnover. This is likely due to the fact that  $\beta_{\text{Replacement}}$  is not normalized (by either  $N$  or  $\gamma$ ). Rather, dissimilarity measures should be preferred when the full incidence matrix is available (Roden, Kocsis, Zuschin, & Kiessling, 2018), and  $\beta_{\text{Stan}}$  or an equivalent metric (see Chao & Chiu, 2016; Chao et al., 2012) should be used when it is not available.

Partitioning methods based on only  $\gamma$ - and  $\alpha$ -diversity (i.e., richness-only  $\beta$ -diversity partitioning) intrinsically disregard the species composition of each site, which, depending on the structure of the underlying incidence matrix, may have a disproportionate effect on the accuracy of partitioned measures. This is rather intuitive: for a given combination of  $\gamma$ - and  $\alpha$ -diversity values, one can generate a very large number of different matrices, due to the fact that neither  $\gamma$  nor  $\alpha$  include information on the number of sites in the system. However, even if we fix the number of sites to a given value  $N$  (since, in reality, the number of islands in an archipelago should be known), the number of possible matrices with  $\gamma$  species,  $N$  sites and average species richness per site equal to  $\alpha$  might still be very large. In turn, this means that a given partitioned measure of  $\beta$ -diversity or nestedness may potentially correspond to a broad array of different values of their matrix-wide counterparts. To illustrate this concept,



**FIGURE 3** The range of possible values for NODF and Sørensen multi-site dissimilarity ( $\beta_{Sor}$ ) for five incidence matrices, given the constraints of a dataset's  $\gamma$ -diversity,  $\alpha$ -diversity and number of islands. For a given matrix the coloured rectangle represents the range of possible values, and the same coloured dot represents the observed values for that matrix. In each case, the procedure consisted of attempting a large number of random swaps (10,000) between species presences/absences within sites, retaining only those swaps that moved the matrix structure in the desired direction (i.e., towards the extremes of possible NODF and  $\beta_{Sor}$  values). Five incidence matrices were randomly selected from the full set of 254. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

we chose a random block of 20 matrices from the set of 254 incidence matrices used in the main analysis, and we used a simple procedure to explore how much, for each matrix, we could modify the matrix structure towards either higher or lower  $\beta$ -diversity and nestedness relative to the observed values, without altering  $\gamma$ ,  $\alpha$  or  $N$  (see Supporting Information Appendix S3 for details). For clarity, results for a subset of five matrices are reported in Figure 3, whilst we provide separate plots for each of the 20 matrices in Figure S2 in Supporting Information Appendix S3. As expected, for a given matrix, the same  $\gamma$ ,  $\alpha$  and  $N$  can result in a wide range of internal matrix structure, especially in terms of nestedness. In turn, this makes it very difficult to draw parallels between the matrix-wide and the partitioned concepts of turnover and nestedness, as the latter might span a very large spectrum of cases that can only be finely discriminated using the former.

The results of our main analyses were largely consistent for the different subsets of datasets, although there were a few subsets ( $\alpha > 25$ ,  $\gamma < 20$  and  $\gamma > 77$ ) for which the correlation between  $\beta_{Nestedness}$  and  $\beta_{Sne}$  was positive and significant. In datasets with low  $\gamma$ , the richness differences between islands are likely constrained and thus  $\beta_{Nestedness}$  and  $\beta_{Sne}$  are both restricted to low values, which could explain the positive correlation between  $\beta_{Nestedness}$  and  $\beta_{Sne}$  for the low  $\gamma$  subset. The reason for the positive correlations observed in the high  $\gamma$  and high  $\alpha$  subsets is unclear, but may point towards a joint dependency between the metrics and  $\gamma$ . That being said, it should be noted that, whilst the correlations were

significant, the coefficients were relatively low (i.e., .23, .24 and .44) and thus our ability to make conclusions based on these results is limited.

The results of the present study clearly illustrate the issues with using richness-only  $\beta$ -diversity partitioning to measure species turnover and nestedness, and we recommend that, if using this approach, more accurate definitions are adopted for these terms in future studies. Readers are directed to Ulrich et al. (2009), Baselga (2012) and Chao and Chiu (2016) for discussion of other nestedness and  $\beta$ -diversity metrics. More generally, the results of this study also highlight the benefits of the deposition of datasets from published studies in data archives.








## ACKNOWLEDGMENTS

José María Fernández-Palacios and Juliano Cabral kindly reviewed an earlier version of the paper. Brian McGill, Petr Keil and two anonymous referees provided useful comments that greatly improved the paper.

## DATA ACCESSIBILITY

The incidence matrices and the R code used to run the analyses are provided in a GitHub repository (<https://github.com/txm676/Partitioning-additive-beta>). The repository has been archived on the Zenodo research data repository (<https://doi.org/10.5281/zenodo.2595322>).

## ORCID

Thomas J. Matthews  <https://orcid.org/0000-0002-7624-244X>  
 Werner Ulrich  <https://orcid.org/0000-0002-8715-6619>  
 Yasuhiro Kubota  <https://orcid.org/0000-0002-5723-4962>  
 Konstantinos Proios  <https://orcid.org/0000-0001-7545-5892>  
 Kostas A. Triantis  <https://orcid.org/0000-0003-2737-8890>  
 Robert J. Whittaker  <http://orcid.org/0000-0001-7775-3383>  
 Giovanni Strona  <https://orcid.org/0000-0003-2294-4013>

## REFERENCES

- Almeida-Neto, M., Frensel, D. M. B., & Ulrich, W. (2012). Rethinking the relationship between nestedness and beta diversity: A comment on Baselga (2010). *Global Ecology and Biogeography*, 21, 772–777. <https://doi.org/10.1111/j.1466-8238.2011.00709.x>
- Almeida-Neto, M., Guimarães, P., Guimarães Jr, P. R., Loyola, R. D., & Ulrich, W. (2008). A consistent metric for nestedness analysis in ecological systems: Reconciling concept and measurement. *Oikos*, 117, 1227 – 1239. <https://doi.org/10.1111/j.0030-1299.2008.16644.x>
- Anderson, M. J., Crist, T. O., Chase, J. M., Vellend, M., Inouye, B. D., Freestone, A. L., ... Swenson, N.G. (2011). Navigating the multiple meanings of  $\beta$  diversity: A roadmap for the practicing ecologist. *Ecology Letters*, 14, 19–28.

- Atmar, W., & Patterson, B. D. (1995). *The nestedness temperature calculator: A Visual Basic program*. Chicago, IL: AICS Research. Retrieved from <http://aics-research.com/nestedness/tempcalc.html>
- Baselga, A. (2010a). Multiplicative partition of true diversity yields independent alpha and beta components; additive partition does not. *Ecology*, *91*, 1974–1981.
- Baselga, A. (2010b). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, *19*, 134–143.
- Baselga, A. (2012). The relationship between species replacement, dissimilarity derived from nestedness, and nestedness. *Global Ecology and Biogeography*, *21*, 1223–1232. <https://doi.org/10.1111/j.1466-8238.2011.00756.x>
- Baselga, A., & Orme, C. D. L. (2012). Betapart: An R package for the study of beta diversity. *Methods in Ecology and Evolution*, *3*, 808–812.
- Cabral, J. S., Weigelt, P., Kissling, W. D., & Krefl, H. (2014). Biogeographic, climatic and spatial drivers differentially affect  $\alpha$ -,  $\beta$ - and  $\gamma$ -diversities on oceanic archipelagos. *Proceedings of the Royal Society B: Biological Sciences*, *281*, 20133246.
- Carvalho, J. C., Cardoso, P., & Gomes, P. (2012). Determining the relative roles of species replacement and species richness differences in generating beta-diversity patterns. *Global Ecology and Biogeography*, *21*, 760–771. <https://doi.org/10.1111/j.1466-8238.2011.00694.x>
- Chao, A., & Chiu, C.-H. (2016). Bridging the variance and diversity decomposition approaches to beta diversity via similarity and differentiation measures. *Methods in Ecology and Evolution*, *7*, 919–928. <https://doi.org/10.1111/2041-210X.12551>
- Chao, A., Chiu, C.-H., & Hsieh, T. C. (2012). Proposing a resolution to debates on diversity partitioning. *Ecology*, *93*, 2037–2051. <https://doi.org/10.1890/11-1817.1>
- Chiarucci, A., Bacaro, G., Arévalo, J. R., Delgado, J. D., & Fernández-Palacios, J. M. (2010). Additive partitioning as a tool for investigating the flora diversity in oceanic archipelagos. *Perspectives in Plant Ecology, Evolution and Systematics*, *12*, 83 – 91. <https://doi.org/10.1016/j.ppees.2010.01.001>
- Cowie, R. H. (1995). Variation in species diversity and shell shape in Hawaiian Land Snails: In situ speciation and ecological relationships. *Evolution*, *49*, 1191–1202. <https://doi.org/10.1111/j.1558-5646.1995.tb04446.x>
- Crist, T. O., & Veech, J. A. (2006). Additive partitioning of rarefaction curves and species–area relationships: Unifying  $\alpha$ -,  $\beta$ - and  $\gamma$ -diversity with sample size and habitat area. *Ecology Letters*, *9*, 923–932. <https://doi.org/10.1111/j.1461-0248.2006.00941.x>
- Crist, T. O., Veech, J. A., Gering, J. C., & Summerville, K. S. (2003). Partitioning species diversity across landscapes and regions: A hierarchical analysis of  $\alpha$ ,  $\beta$ , and  $\gamma$  diversity. *The American Naturalist*, *162*, 734–743. <https://doi.org/10.1086/378901>
- Jost, L. (2007). Partitioning diversity into independent alpha and beta components. *Ecology*, *88*, 2427–2439. <https://doi.org/10.1890/06-1736.1>
- Koleff, P., Gaston, K. J., & Lennon, J. J. (2003). Measuring beta diversity for presence-absence data. *Journal of Animal Ecology*, *72*, 367–382. <https://doi.org/10.1046/j.1365-2656.2003.00710.x>
- Lande, R. (1996). Statistics and partitioning of species diversity, and similarity among multiple communities. *Oikos*, *76*, 5–13. <https://doi.org/10.2307/3545743>
- Legendre, P., & De Cáceres, M. (2013). Beta diversity as the variance of community data: Dissimilarity coefficients and partitioning. *Ecology Letters*, *16*, 951–963. <https://doi.org/10.1111/ele.12141>
- Legendre, P., & Legendre, L. (1983). *Numerical ecology*. Amsterdam, the Netherlands: Elsevier.
- MacArthur, R., Recher, H., & Cody, M. (1966). On the relation between habitat selection and species diversity. *The American Naturalist*, *100*, 319–332. <https://doi.org/10.1086/282425>
- Matthews, T. J. (2015). Analysing and modelling the impact of habitat fragmentation on species diversity: A macroecological perspective. *Frontiers of Biogeography*, *7*, 60–68. <https://doi.org/10.21425/F5FBG25942>
- Matthews, T. J., Cottee-Jones, H. E. W., & Whittaker, R. J. (2015). Quantifying and interpreting nestedness in habitat islands: A synthetic analysis of multiple datasets. *Diversity and Distributions*, *21*, 392–404. <https://doi.org/10.1111/ddi.12298>
- Matthews, T. J., Guilhaumon, F., Triantis, K. A., Borregaard, M. K., & Whittaker, R. J. (2016). On the form of species–area relationships in habitat islands and true islands. *Global Ecology and Biogeography*, *25*, 847–858.
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., ... Wagner, H. (2017). *Vegan: Community ecology package*. R package version 2.4-5. Retrieved from <https://cran.r-project.org/web/packages/vegan/index.html>
- Patterson, B. D., & Atmar, W. (1986). Nested subsets and the structure of insular mammalian faunas and archipelagos. *Biological Journal of the Linnean Society*, *28*, 65–82. <https://doi.org/10.1111/j.1095-8312.1986.tb01749.x>
- R Core Team. (2017). R: A language and environment for statistical computing (Version 3.5.2). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Roden, V. J., Kocsis, Á. T., Zuschin, M., & Kiessling, W. (2018). Reliable estimates of beta diversity with incomplete sampling. *Ecology*, *99*, 1051–1062. <https://doi.org/10.1002/ecy.2201>
- Rosenzweig, M. L. (1995). *Species diversity in space and time*. UK: Cambridge University Press.
- Sfenthourakis, S., & Panitsa, M. (2012). From plots to islands: Species diversity at different scales. *Journal of Biogeography*, *39*, 750–759. <https://doi.org/10.1111/j.1365-2699.2011.02639.x>
- Strona, G., Ulrich, W., & Gotelli, N. J. (2017). Bi-dimensional null model analysis of presence-absence binary matrices. *Ecology*, *99*, 103–115. <https://doi.org/10.1002/ecy.2043>
- Triantis, K. A., Guilhaumon, F., & Whittaker, R. J. (2012). The island species–area relationship: Biology and statistics. *Journal of Biogeography*, *39*, 215–231. <https://doi.org/10.1111/j.1365-2699.2011.02652.x>
- Tuomisto, H. (2010). A diversity of beta diversities: Straightening up a concept gone awry. Part 1. Defining beta diversity as a function of alpha and gamma diversity. *Ecography*, *33*, 2–22. <https://doi.org/10.1111/j.1600-0587.2009.05880.x>
- Ulrich, W., Almeida-Neto, M., & Gotelli, N. J. (2009). A consumer's guide to nestedness analysis. *Oikos*, *118*, 3–17. <https://doi.org/10.1111/j.1600-0706.2008.17053.x>
- Veech, J. A., & Crist, T. O. (2010). Toward a unified view of diversity partitioning. *Ecology*, *91*, 1988–1992. <https://doi.org/10.1890/09-1140.1>
- Veech, J. A., Summerville, K. S., Crist, T. O., & Gering, J. C. (2002). The additive partitioning of species diversity: Recent revival of an old idea. *Oikos*, *99*, 3–9. <https://doi.org/10.1034/j.1600-0706.2002.990101.x>
- Whittaker, R. H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecological Monographs*, *30*, 279–338. <https://doi.org/10.2307/1943563>
- Whittaker, R. H. (1965). Dominance and diversity in land plant communities. *Science*, *147*, 250–260.
- Whittaker, R. J., & Fernández-Palacios, J. M. (2007). *Island biogeography: Ecology, evolution, and conservation* (2nd ed.). UK: Oxford University Press.
- Zhang, Q., Hou, X., Li, F. Y., Niu, J., Zhou, Y., Ding, Y., ... Kang, S. (2014). Alpha, beta and gamma diversity differ in response to precipitation in the inner Mongolia grassland. *PLoS ONE*, *9*, e93518. <https://doi.org/10.1371/journal.pone.0093518>



## BIOSKETCH

**TOM MATTHEWS** is a macroecologist and biogeographer at the University of Birmingham, UK. He is interested in the application of macroecological methods to global environmental change questions, and his previous work has focused on the impacts of habitat fragmentation and the form of the species–area relationship in fragmented landscapes.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Matthews TJ, Aspin TWH, Ulrich W, et al. Can additive beta diversity be reliably partitioned into nestedness and turnover components? *Global Ecol Biogeogr.* 2019;28:1146–1154. <https://doi.org/10.1111/geb.12921>